

Document made available under the Patent Cooperation Treaty (PCT)

International application number: PCT/US05/005309

International filing date: 18 February 2005 (18.02.2005)

Document type: Certified copy of priority document


Document details: Country/Office: US
Number: 60/548,403
Filing date: 27 February 2004 (27.02.2004)

Date of receipt at the International Bureau: 17 March 2005 (17.03.2005)

Remark: Priority document submitted or transmitted to the International Bureau in compliance with Rule 17.1(a) or (b)



World Intellectual Property Organization (WIPO) - Geneva, Switzerland
Organisation Mondiale de la Propriété Intellectuelle (OMPI) - Genève, Suisse



THE UNITED STATES OF AMERICA

TO ALL TO WHOM THESE PRESENTS SHALL COME:

UNITED STATES DEPARTMENT OF COMMERCE

United States Patent and Trademark Office

March 08, 2005

THIS IS TO CERTIFY THAT ANNEXED HERETO IS A TRUE COPY FROM THE RECORDS OF THE UNITED STATES PATENT AND TRADEMARK OFFICE OF THOSE PAPERS OF THE BELOW IDENTIFIED PATENT APPLICATION THAT MET THE REQUIREMENTS TO BE GRANTED A FILING DATE.

APPLICATION NUMBER: 60/548,403

FILING DATE: February 27, 2004

RELATED PCT APPLICATION NUMBER: PCT/US05/05309



Certified by


Under Secretary of Commerce
for Intellectual Property
and Director of the United States
Patent and Trademark Office

PROVISIONAL APPLICATION COVER SHEET

This is a request for filing a PROVISIONAL APPLICATION under 35 CFR 1.53 (c).

Docket Number		62644	Type a plus sign (+) inside this box	+
INVENTOR(s)/APPLICANT(s)				
LAST NAME	FIRST NAME	MIDDLE INITIAL	RESIDENCE (CITY AND EITHER STATE OR FOREIGN COUNTRY)	
Frank Donate Shields Li	Timothy Felipe Jefry Kai	C. A. E.	5001 Foxpoint Circle, Midland, Michigan 48642 4608 Lund, Midland, Michigan 48642 1066 Fulton Road, San Marcos, California 92069 1434 Sundance Avenue, San Marcos, California 92078	
<input type="checkbox"/> Additional inventors are being named on separately numbered sheets attached hereto				
TITLE OF THE INVENTION (280 characters max)				
METHOD FOR THE EXTRACTION OF INTRACELLULAR PROTEINS FROM A FERMENTATION BROTH				
CORRESPONDENCE ADDRESS				
Dragan J. Karadzic THE DOW CHEMICAL COMPANY Intellectual Property P. O. Box 1967 Midland, MI 48641-1967 U.S.A.			Telephone: 979-238-2845 Fax: 979-238-0878	
ENCLOSED APPLICATION PARTS (check all that apply)				
<input checked="" type="checkbox"/> Specification		Number of Pages	28	<input type="checkbox"/> Small Entity Statement
<input type="checkbox"/> Drawing(s)		Number of Sheets	_____	<input type="checkbox"/> Other (specify) _____
METHOD OF PAYMENT (check one)				
<input type="checkbox"/> A check or money order is enclosed to cover the Provisional filing fees			PROVISIONAL FILING FEE AMOUNT	\$160.00
<input checked="" type="checkbox"/> The Commissioner is hereby authorized to charge filing fees and credit Deposit Account Number: No. 04-1512				
The invention was made by an agency of the United States Government or under a contract with an agency of the United States Government				
<input checked="" type="checkbox"/> No				
<input type="checkbox"/> Yes, the name of the U.S. Government agency and the Government contract number are: _____				

Respectfully submitted,



Dragan J. Karadzic
Registration No.: 28,024
Phone No.: (979) 238-2845

DJK/mfg

EXPRESS MAIL MAILING NO.: EV370690583US
DATE OF DEPOSIT: February 27, 2004

PROVISIONAL APPLICATION FILING ONLY

METHOD FOR THE EXTRACTION OF INTRACELLULAR PROTEINS FROM A FERMENTATION BROTH

Background of the Invention

5 The present invention relates to a process for the extraction of intracellular proteins including enzymes and therapeutic proteins from fermentation broth using an organic solvent.

10 Enzymes are highly efficient protein catalysts which are involved in almost every biological reaction. The enzymes are grouped in into six major classes on the basis of the type of reaction catalyzed; i.e., Oxidoreductase, Transferase, Hydrolase, Lyase, Isomerase and Ligase. Enzymes find use in chemical analysis, clinical diagnosis and a broad range of industrial applications. Enzymes may be extracted from any living organism but most are obtained by the fermentation of micro-organisms. Industrial
15 preparation of industrial enzymes aims for economy, effectiveness and safety.

 Therapeutic proteins are proteins with specific biological activity that make them effective as pharmaceutical agents or drugs for treatment of disease, or as adjuncts to therapy used in combination with a drug or mixture of drugs. In certain cases,
20 therapeutic proteins may be produced via bacterial fermentation using the methods of genetic engineering to cause the host microbe to produce a specific protein or mixture of proteins having the desired activity. Often, these recombinant proteins are produced within the cell of the organism and must be recovered from the cell after harvesting the broth. Because therapeutic proteins are used as drugs or in combination with drugs, the
25 purity of the isolated protein or protein mixture is a critical factor in their manufacture. This is especially important for proteins that must be injected into the bloodstream to effectively treat disease.

 Methods used to recover proteins from fermentation broths can essentially be
30 categorized in two basic methods:

 (i) when the host cells are able to secrete proteins outside the cells into a growth medium (extracellular proteins), the media in which the cells grow is collected and the proteins are harvested from the liquid phase, sometimes as the cells continue to grow. This is desirable as it provides the protein in a medium that is not loaded with other
35 cellular components like DNA, host cell proteins, etc. This is typically done in mammalian cell systems but there have been recent advances in prokaryotic cell

systems like *E. coli*. The recovery of the desired protein from the media in which the cell grows involves filtration, chromatography, etc.; and

(ii) more commonly in bacterial cell lines like *E. coli* and *P. fluorescens*, the protein is produced and sequestered inside the cell (intracellular protein). In order to
5 recover the intracellular protein it is necessary first to get it out of the cell. Cellular disruption techniques like microfluidization, osmotic shock, heating, pH adjustment work well often but the difficulty in isolating the desired protein from the rest of the cellular components becomes a challenge as once the cell's integrity is disrupted all of the cell's proteins, DNA and other biomolecules, are spilled into the solution that
10 contains the targeted protein. Additionally, when bacterial cells over express a protein/enzyme, the vast amounts they are able to produce have difficulty adopting the correct conformation that imparts their special activity. These large amounts of protein begin to accumulate and congregate together forming insoluble particles within the cell that are called inclusion bodies. Often these proteins need to be unfolded into a
15 linear amino acid sequence and then refolded to yield an active protein. Techniques for recovery from these complex biological soups require solid/liquid separations, filtrations, precipitations, chromatography, etc.

Some intracellular enzymes are used commercially without isolation and
20 purification but the majority of commercial enzymes and therapeutic proteins are either produced extracellularly by the microbe and must be recovered from the liquid phase, or they are produced intracellularly by the microbe and must be recovered from the cells and further processed. For recovery of intracellular proteins at an industrial scale, solids/liquid separation is generally required for separation of cell mass, the removal of
25 cell debris after cell breakage and the collection of precipitates. This can be done by filtration or centrifugation. In general, filtration is used to remove unwanted cells or cell debris whereas centrifugation is a preferred method for the collection of required solid material.

30 Various methods for recovery and/or purification of proteins are described in the literature. These methods include cell disruption, microfiltration, ultrafiltration, various forms of chromatography and ion exchange, as well as aqueous two-phase extraction (also called aqueous biphasic partitioning) using water-soluble polymers (including polyglycols and nonionic surfactants). See, for example, M. P. Deutscher, Ed., "Guide
35 to Protein Purification," Methods in Enzymology," Vol. 182, Academic Press, San Diego,(1990); H. B. Blanch and D. S. Clark, "Biochemical Engineering," Marcel

Dekker, New York, 1997, pp. 474-482; P. A. Belter, E. L. Cussler and W. Hu, "Bioseparations: Downstream Processing for Biotechnology," Wiley, New York, 1988, Chapter 5, 55. 99-143; and M.R. Ladisch, "Bioseparation Engineering," Wiley Interscience, New York, 2001.

5

Aqueous two-phase extraction has been widely used for the separation and concentration of proteins and nucleic acids. The two-phase aqueous systems are generally made up of (i) two immiscible polymer components, both water-soluble, such as polyethylene glycol and dextran; or (i) a single polymer component, such as polyethylene glycol, and aqueous salt solution; or (iii) water-miscible organic solvent, such as ethanol, and aqueous salt solution; or (iv) a non-ionic detergent and hydrophilic polymers, such as polyethylene glycol and dextran. See, for example, A. Louwrier, "Model Phase Separations of Proteins Using Aqueous/Ethanol Components", Biotechnology Techniques, Vol. 12, No. 6, pp. 363-365 (1998); A. Louwrier, "Model Isolations of Nucleic Acids from Prokaryotic Sources Using an Inorganic/Aqueous Biphasic System", Biotechnology Techniques, 13, pp 329-330 (1999); A. Louwrier, "Nucleic Acid Removal from *Taq* Polymerase Preparations Using an Aqueous/Organic Biphasic System", Biotechnology Techniques, 27, pp 444-445 (1999); Ulf Sivars et al, "Mechanism of Phase Behavior and Protein Partitioning in Detergent/Polymer Aqueous Two-Phase Systems for Purification of Integral Membrane Proteins", Biochemica and Biophysica Acta, 1474, pp 133-146 (2000); and Jorge Lorwin et al, "Oxidative Renaturation of Hen Egg-White Lysosyme in Polyethylene-Salt Aqueous Two-Phase Systems", Biotechnology and Bioengineering, Vol.65, No.4, pp. 437-446 (1999),

25

British Patent No. 2,333,526A discloses a process for extracting a nucleic acid from biochemical material using a biphasic system made of (i) a water miscible organic solvent which is preferably short chain alcohol such as methanol; and (ii) water, in combination with a partitioning agent. The two solvents, the water miscible organic solvent and water, are normally 100% miscible. When a partitioning agent, which is preferably an inorganic salt such as a phosphate, sulphate or carbonate, is added to the water component a biphasic system is formed. The aqueous phase contains nucleic acid and the organic phase contains the majority of proteins. The organic and aqueous phases are then separated by decantation or centrifugation.

35

International Patent Publication No. WO 00/12537 discloses a process for the preparation of biologically active somatotropin from inclusion bodies of a recombinant host cell containing an inactive form of the somatotropin protein involving the steps of: (a) contacting the inclusion bodies with an aqueous alcohol solution, particularly an aqueous n-propyl or isopropyl alcohol, at an alkaline pH to solubilize the protein; and (b) bringing the solubilized protein into contact with a mild oxidising agent to refold and form intramolecular disulfide bond between cysteine residues of the protein.

Hans-Olof Johansson et al, "Thermoseparating Water/Polymer System: A Novel One-Polymer Aqueous Two-Phase System for Protein Purification", Biotechnology and Bioengineering, Vol. 66, No. 4, pp. 247-257 (1999), describes the partitioning and separation of proteins using an aqueous two-phase system composed of only one polymer in water solution. The polymer is a linear random copolymer composed of ethylene oxide and propylene oxide groups which has been hydrophobically modified with myristyl groups at both ends. The polymer thermoseparates in water with cloud point at 14 °C.

The known processes for the extraction/purification of intracellular proteins are not efficient and require disruption of cellular walls, folding and refolding of the proteins. Because the cell wall must be disrupted, the contents of the cell are released and mixed with the desired protein. The isolation of the active protein from this soup of impurities is difficult and expensive due to the requirement for numerous processing steps needed to deal with the variety and quantity of impurities. Thus, there exists a need for a more effective and efficient method for extraction of intracellular proteins. An improved method for the extraction of a protein from a fermentation broth that does not disrupt cell walls and does not require folding and refolding of enzymes has now been discovered. This new method uses certain oxygenated organic solvents that are water miscible or partially water miscible.

Summary of the Invention

The invention involves the use of water miscible alcohols, water miscible glycol ethers, or partially water miscible glycol ethers to extract intracellular protein from whole fermentation broth. In the context of the present invention, the term whole fermentation broth means broth that has not been filtered to remove the biomass including cells. Water miscible alcohols and water miscible glycol ethers are

completely miscible with water in all proportions in the temperature range from 20 C to 80 C. All other alcohols are partially water miscible alcohols and all other glycol ethers are partially water miscible glycol ethers.

5 In one aspect, the present invention concerns a method for extracting an intracellular protein from a fermentation broth comprising the steps of:

- 10 (a) intermixing a sufficient quantity of a water miscible alcohol or glycol ether with an aqueous fermentation broth at a temperature at which the alcohol or glycol ether forms a single aqueous solution comprising a protein, the water miscible alcohol or glycol ether, and water;
- 15 (b) separating the aqueous phase comprising the protein, the water miscible alcohol or glycol ether, and water formed in step (a) from solid biomass impurities; and, optionally,
- (c) recovering the protein from the aqueous phase obtained in step (b) by any conventional protein recovery method.

20 In another aspect, the present invention concerns a method for extracting an intracellular protein from a fermentation broth comprising the steps of:

- 25 (a) intermixing a sufficient quantity of a partially water miscible glycol ether with an aqueous fermentation broth at a temperature at which the glycol ether forms a single aqueous solution comprising a protein, the partially water miscible glycol ether, and water;
- 30 (b) separating the aqueous phase comprising the protein, the partially water miscible glycol ether, and water formed in step (a) from solid biomass impurities; and, optionally,
- (c) recovering the protein from the aqueous phase obtained in step (b) by any conventional protein recovery method.

35 In still another aspect, the present invention concerns a method for extracting an intracellular protein from a fermentation broth comprising the steps of:

- 5 (a) intermixing a sufficient quantity of a partially water miscible glycol ether with an aqueous fermentation broth at a temperature where the glycol ether forms an aqueous phase comprising a protein, partially water miscible glycol ether, and water; and an organic phase comprised mainly of partially miscible glycol ether;
- 10 (b) separating the liquid organic phase formed in step (a) from the aqueous phase,
- (d) separating the aqueous phase obtained in step (b) from solid biomass impurities; and, optionally,
- 15 (d) recovering the protein from the aqueous phase obtained in step (c) by any conventional protein recovery method.

Detailed Description of the Invention

20 The method of the present invention is useful for the extraction of proteins from an aqueous fermentation broth. The method of the present invention is particularly useful for the extraction of proteins expressed as inclusion bodies within the cell of the organism. In the context of the present invention, the term "inclusion bodies" refers to cytoplasmic aggregates containing heterologous proteins expressed in a transformed host cell, which can be recovered by separating from the cytoplasm.

25 For example, the present invention is suitable for, but not limited to, the extraction of an amylase enzyme from a whole fermentation broth. Amylases are one of the most commonly used classes of industrial enzymes. Amylases find applications in starch processing, baking, brewing, alcohol production, textile and other industries.

30 In this example, α -amylase enzyme is extracted from the fermentation broth in a single liquid phase using an aqueous solution comprising water and an alcohol, water miscible glycol ether or a partially water miscible glycol ether without disrupting cells integrity and without pretreatment steps specifically designed to lyse the cells. The extracted enzyme maintains most of its activity.

35

Non-limiting examples of alcohols that are useful in the practice of the present invention are methanol, ethanol, 2-propanol (IPA), and 2-methyl-2-propanol. The alcohol 2-propanol is a preferred alcohol.

5 A mixture of two or more alcohols can also be used in the method of the present invention to tailor the solvent for a specific protein and optimize performance.

Glycol ethers that are useful in the practice of the present invention are water miscible or partially water miscible. The glycol ethers exhibit both hydrophobic and hydrophilic characteristics because of the presence of hydrophobic alkyl groups and hydrophilic oxygen-containing functional groups (hydroxyl groups and ether linkages).
10 Some of glycol ethers also exhibit relatively low aquatic and mammalian toxicity compared with other organic solvents.

15 Many glycol ethers exhibit inverse solubility in water such that the solubility in water at 100 °C is at least 1 weight percent less than the solubility in water at -5 °C. This inverse solubility behavior can be attributed to temperature-sensitive hydrogen bonding. It is known that many glycol ethers exhibit a lower critical solution temperature (LCST) below which they are completely miscible with water. At
20 temperatures below the LCST, the glycol ether is able to form hydrogen bonds with water and this attractive interaction leads to complete miscibility. At temperatures above the LCST, hydrogen bonding is disrupted by increasing thermal energy and hydrophobic interactions between the glycol ether and water begin to dominate. This results in partial miscibility and a decrease in the solubility of the glycol ether in water
25 with increasing temperature (termed inverse solubility). Depending on the particular glycol ether, the LCST can be as low as -10 °C or as high as 100 °C. The glycol ethers, both water soluble and partially water soluble are well known in the art and various methods for their preparation are described in the literature and practiced commercially.

30 Non-limiting examples of water miscible glycol ethers that are useful in the practice of the present invention and are completely miscible with water in the temperature range of from about 20 °C to about 80 °C are: ethylene glycol n-propyl ether (EP), propylene glycol ethyl ether (PE), propylene glycol methyl ether (PM), diethylene glycol n-butyl ether (DB), diethylene glycol ethyl ether (DE), diethylene
35 glycol methyl ether (DM), triethylene glycol n-butyl ether (TB), triethylene glycol n-

pentyl ether (TPent), triethylene glycol ethyl ether (TE), triethylene glycol methyl ether (TM), and diethylene glycol dimethyl ether (DGDME).

Non-limiting examples of partially water miscible glycol ethers that are useful
 5 in the practice of the present invention and are miscible with water at the temperature of
 about 20 °C but become only partially miscible with water as the water-ethylene glycol
 solution is heated are: ethylene glycol n-butyl ether (EB), ethylene glycol iso-butyl ether
 (EiB), propylene glycol n-propyl ether (PnP), dipropylene glycol ethyl ether (DPE),
 dipropylene glycol iso-propyl ether (DPiP), Diethylene glycol 2-methylbutyl ether
 10 (D2MB), diethylene glycol n-pentyl ether (DPent), triethylene glycol n-heptyl ether
 (THept), triethylene glycol n-hexyl ether (THex), diethylene glycol ethyl ether acetate
 (DEA), and diethylene glycol diethyl ether (DGDEE).

Non-limiting examples of partially water miscible glycol ethers that are useful
 15 in the practice of the present invention that form a separate phase with water at about 20
 °C and separate further upon heating are: ethylene glycol 2-methylbutyl ether (E2MB),
 ethylene glycol n-hexyl ether (EHex), ethylene glycol n-pentyl ether (EPent), propylene
 glycol n-butyl ether (PnB), propylene glycol tert-butyl ether (PtB), propylene glycol iso-
 propyl ether (PiP), dipropylene glycol n-butyl ether (DPnB), dipropylene glycol n-
 20 propyl ether (DPnP), diethylene glycol n-hexyl ether (DHex), tripropylene glycol n-
 butyl ether (TPnB), tripropylene glycol n-propyl ether (TPnP), ethylene glycol ethyl
 ether acetate (EEA), ethylene glycol n-butyl ether acetate (EBA), diethylene glycol n-
 butyl ether acetate (DBA), propylene glycol methyl ether acetate (PMA), ethylene
 glycol diethyl ether (EGDEE), ethylene glycol dibutyl ether (EGDBE), diethylene
 25 glycol dibutyl ether (DGDBE), and dipropylene glycol dimethyl ether (DMM).

A mixture of two or more glycol ethers can also be used in the method of the
 present invention to tailor the solvent for a specific protein and optimize performance.

30 According to the present invention, a protein is extracted from a whole aqueous
 fermentation broth using an alcohol, water miscible glycol ether or a partially water
 miscible glycol ether. The exact nature of the action of the alcohol, water miscible
 glycol ether or partially water miscible glycol ether in the method of the present
 invention is not known, but it is believed that the invention facilitates movement of the
 35 protein present in the biomass solids through the cell wall into an aqueous solution on
 addition of the alcohol, water miscible glycol ether or partially water miscible glycol

ether. The cell wall remains intact and the majority of alpha-amylase enzyme activity is retained.

5 We have discovered that the alpha-amylase enzyme will not transfer from the resulting aqueous solution into a glycol ether phase when a partially water miscible glycol ether is used. This provides for the possibility of further purifying and concentrating the aqueous enzyme obtained in the whole broth extraction by the present method via liquid-liquid extraction of the more lipophilic impurities and some of the water from the aqueous phase into a glycol ether phase.

10

In certain cases, it is advantageous to heat the fermentation broth followed by the addition of a small amount of ethylenediamine-tetraacetic acid (EDTA) or another chelating compound prior to the extraction of a protein to remove metal ions from the cell walls to increase the permeability of the cell walls.

15

Conveniently, the method of the present invention is conducted at a temperature of from about 1 °C to about 100 °C, preferably from about 20 °C to about 40 °C.

20 The biomass impurities present in the aqueous protein solution can be removed via any known method such as, for example, via centrifugation or microfiltration. High speed batch centrifugation can be used for removal of biomass impurities.

25 When a partially water miscible glycol ether is used in the process of the present invention and this results in formation of an organic phase comprised mainly of the partially water miscible glycol ether, the glycol ether can be recovered from the liquid organic phase and recycled into the method. The recovery and recycling of the glycol ether should be achieved with little loss of the glycol ether in order to achieve good method economics. Methods that can be used to recover the partially water miscible glycol ether from the organic phase include distillation, evaporation, and
30 chromatography. Distillation is a preferred method.

Also, if desired, the alcohol and water miscible glycol ether can be recovered from the single aqueous phase comprising the protein, alcohol or glycol ether, as can the glycol ether dissolved in the aqueous phase when a partially miscible glycol ether is
35 used. Distillation is a preferred method for recovery of the alcohol. The recovery of dissolved partially water miscible glycol ether can be facilitated by heating the aqueous

solution to reduce the saturation amount of glycol ether present in the aqueous phase. Recovery partially water miscible glycol ether from the aqueous phase may be accomplished by steam stripping provided that the partially water miscible glycol ether is sufficiently volatile or hydrophobic. In addition, water miscible or partially water miscible glycol ethers may be recovered from aqueous solutions via liquid-liquid
5 extraction using hydrophobic organic solvents such as 2-ethyl hexanol.

If desired, the protein product can be concentrated by any known concentration method such as, for example, by extraction of water into a water-lean glycol ether (that is, one that is not saturated with water), or by evaporation such as in a wiped-film
10 evaporator.

The temperature at which the intermixing of the organic oxygenated solvent with the aqueous fermentation broth in step (a) of the method of the present invention is not critical and can conveniently be from about 1 °C to about 90 °C, preferably from about 20 °C to about 40 °C. The temperature used is dependent upon the particular glycol ether used and the particular protein.
15

The method of the present invention is advantageously carried out at atmospheric pressure, although higher and lower pressures may be used in certain cases.
20

A person skilled in the art may readily select the amount of an alcohol or glycol ether which may be used. Generally speaking, a sufficient amount of the alcohol or glycol ether must be used to obtain substantial recovery of the desired protein from the aqueous fermentation broth. This can be readily determined by experimentation.
25

The method of the present invention can be carried out in a batch operation or continuously, and may be conducted in any conventional extraction equipment.

If desired, additional equipment can be used in the method of the present invention such as additional extraction, distillation or evaporation equipment for recovery of dissolved glycol ether from the aqueous phase and the organic phase, and for recovery and recycle of the hydrophobic co-solvent, when used. Such additional equipment and its use in liquid-liquid extraction methods are well known in the art. A
30 person of an ordinary skill in the art would use such additional equipment or
35 combination thereof in the method of the present invention in a manner known for use

of such equipment in conventional liquid-liquid extraction methods. The use of such additional equipment or combination thereof will depend on many factors, such as, for example, the nature of the protein, the nature of other compounds present in the fermentation broth, the nature of the glycol ether used, the use of the hydrophobic organic solvent, the costs associated with the use of glycol ether and/or hydrophobic solvent, the costs associated with the use of additional equipment, and the overall economics of the entire process.

Examples

The invention is demonstrated by the example of extraction of alpha-amylase enzyme from whole fermentation broth. The enzyme activity after extraction by the method of the present invention is determined by one of two different colorimetric methods. The first method is modeled after procedures given in experiments 10.1 and 10.8 described in David T. Plummer's book, An Introduction to Practical Biochemistry, McGraw-Hill, New York, 1971, where a salivary alpha-amylase enzyme is used instead. This method detects the sugar maltose produced when the alpha-amylase enzyme is used to hydrolyze the alpha-1 \rightarrow 4 links of a starch sample. A spectrophotometer is used to measure the absorbance of the resulting maltose solution at 540 nm. The second method relies on the detection of 4-nitrophenol liberated when the alpha-amylase enzyme is used to hydrolyze a sample of 4-nitrophenyl-alpha-D-hexa-(1 \rightarrow 4)-glucopyranoside substrate. A spectrophotometer is used to measure the absorbance of the resulting 4-nitrophenol solution at 405 nm.

All parts, percentages and ratios herein are by weight unless otherwise indicated.

The invention will be further clarified by a consideration of the following examples which are intended to be purely exemplary of the present invention.

General Test Procedures

Enzyme Activity: The enzyme activity was determined using one of two tests.

In the first test, the activity of the enzyme was assessed using a qualitative Yes/No test. The test was performed on the extracted liquid after extraction using a

starch hydrolysis method with colorimetric detection of the product sugar. The starch substrate was converted to maltose which was then detected using a spectrophotometer to measure UV absorbance. The test procedure is modeled after procedures given in experiments 10.1 and 10.8 of David T. Plummer's book, An Introduction to Practical Biochemistry, McGraw-Hill, New York, 1971, where a salivary alpha-amylase enzyme is used instead.

Preparation of Test Reagents. A phosphate buffer (0.1 M, pH 6.86) was prepared using a commercially available, ready-to-dilute, phosphate buffer salt concentrate, and a 1% NaCl solution was prepared by dissolving the appropriate amount of NaCl in water. A buffered starch substrate was made using the prepared phosphate buffer (0.5% starch in phosphate buffer). Soluble starch (5 g) and a stir bar were placed in a tared beaker which was then placed on a stirring plate. Phosphate buffer (50 ml) was added, and the resulting mixture stirred until a smooth paste was obtained. The paste was added to 500 ml of boiling phosphate buffer and allowed to boil for about one minute. The solution was then cooled to room temperature and diluted to 1 L with the phosphate buffer. A dinitrosalicylate reagent required for the test was prepared as follows: Sodium potassium tartrate (150 g) was dissolved in water (250 ml) in a volumetric flask. The 3,5-dinitrosalicylic acid was placed in a tared beaker along with 2N sodium hydroxide (100 ml) and a magnetic stir bar. The beaker was then placed on a stirred hot-plate and the mixture heated to about 60 °C. The mixture was allowed time to heat and stir until a solution formed. While still hot, the contents of the beaker were placed in a 500-ml volumetric flask that already contained the sodium potassium tartrate solution. The reagent mixture was then diluted to 500 ml.

Test Procedure. Tests were conducted in a series of 50 mL vials, one of which was labeled as the blank. To each of the vials was added 0.5% starch solution (12.5 mL), 0.1 M phosphate buffer (5 mL), and 1% NaCl solution (2.5 mL). Water (2.5 mL) was added to the vial labeled as the blank. A sample of a given aqueous or solvent test solution (2.5 mL) was added to another vial which was then labeled appropriately. This last step was repeated with the remaining test samples, each time using a different vial. The vials were capped and placed in a 37 °C water bath for 20 minutes. The vials were removed from the bath and the enzymatic reaction quenched by the addition of 2N NaOH (2.5 mL) to all of the vials. The dinitrosalicylate reagent (2.5 mL) was added to each of the vials which were then capped and heated for exactly 5 minutes in a boiling water bath. The vials were allowed to cool to room temperature before transferring the

contents to cuvettes. A Hach DR/2010 Spectrophotometer was used to measure the absorbance of each solution at a wavelength of 540 nm. The blank was used to zero the instrument. The test was interpreted to indicate a reduction of activity if the measured absorbance value was lower than the absorbance value measured for a starch sample treated with a stock solution of alpha-amylase. The activity was considered to be fully retained if the absorbance value was the same or higher. The results were reduced to Yes (activity was retained) and No (a significant reduction of activity had occurred). The results obtained are summarized in Tables 2 and 5.

Specific enzyme activity was assessed by measuring the p-nitrophenol liberated by reaction of the enzyme with p-nitrophenyl- α -D-hexa-(1 \rightarrow 4)-glucopyranoside (p-NP substrate).

Preparation of Reagents. A 0.05 M MOPS buffer was prepared by dissolving 11.55g MOPS (3-(N-morpholino) propanesulfonic acid, sodium salt) in 1 L Milli-Q-water (from Millipore Water System) and adjusting the pH to 7 with 1 N HCl. A 6 M urea solution was prepared by dissolving 72 g urea in 200 mL Milli-Q-water. A 5mM p-NP substrate solution was prepared by weighing 111 – 222 mg p-NP into a tared 50-mL plastic centrifuge tube and diluting with 0.05 M MOPS buffer. A standard 10 mM 4-nitrophenol solution was prepared by weighing 140 mg into a tared 100-mL volumetric flask and diluting to volume with 0.05 M MOPS buffer. The pH of the 4-nitrophenol solution was adjusted to 7 with either 1 N HCl or 1 N NaOH. A 1mM 4-nitrophenol calibration standard was prepared by diluting the 10mM solution with 0.05 M MOPS buffer. Alpha-amylase samples and standard solutions were thoroughly stirred prior to sampling. About 1.0 ± 0.1 g alpha-amylase sample or stock solution was weighed in a 5-dram vial and then diluted with 9.0 mL 6 M urea solution. A 1.0 mL aliquot of the resulting urea-enzyme solution was then transferred to a 5-dram vial and diluted with 9.0 mL of the 0.05 MOPS solution.

Test Procedure. Exactly 950 microliters of the 0.05 M MOPS buffer was transferred to a microcuvette and the cuvette placed into a water-heated cuvette heating block at 75 C for several minutes. The cuvette was then removed from the heating block, water wiped off, and placed into the constant temperature reference compartment (75 C) of a Shimadzu UV-161 spectrophotometer. This operation was repeated with another cuvette which was placed in the sample compartment of the spectrophotometer. The absorbance of the cuvette + MOPS buffer was nulled by zeroing the spectrophotometer.

Exactly 50 microliters of the 1mM 4-nitrophenol standard was transferred into the sample cuvette with a micropipette. The solutions were mixed well by gently pipetting up and down several times the liquid in the cuvette, and then by sealing the cuvette and inverting it two times. The absorbance of the solution was then measured three times at 405 nm. Exactly 950 microliters of the 5 mM p-NP substrate solution was transferred to a microcuvette and the cuvette placed into a water-heated cuvette heating block at 75 C for several minutes. The cuvette was removed from the heating block, water wiped off, and placed into the constant temperature sample compartment (75 C) of the spectrophotometer. After several minutes, 50 microliters of the dilute enzyme solution prepared earlier was added to the cuvette. The solutions were mixed well as described before for the 4-nitrophenol reference standards, and then the absorbance of the solution was measured at 405 nm. Enzyme activity concentrations were obtained by comparing the absorbance readings with those obtained with the 1mM p-nitrophenol calibrating standard taking into account dilution factors. The results obtained are summarized in Table 6.

Enzyme concentrations were also established by high pressure liquid chromatography (HPLC) using an Agilent 1100 Series Liquid Chromatograph equipped with an ultraviolet (UV) detector set at 280 nm, and a Dell Ultra Scan P1110 ChemStation. The instrument was fitted with a PLRP-S 5micron, 100A, 250 x 4.6 mm reversed phase column from Polymer Labs. The mobile phase was composed of (A) 0.025 M ammonium acetate (adjusted to pH 9 with NH_4OH) and 10% acetonitrile; and (B) 0.025 M ammonium acetate (adjusted to pH 9 with NH_4OH) and 70% acetonitrile. The analysis used a flow rate of 1 ml/min and a gradient of 100% (A) to 35% (A) in 40 minutes with a 3 minute hold. Injection size was 25 microliters. Samples were prepared by diluting 1.00 g of the aqueous or solvent test solution with 10.00 g HPLC grade water. Calibration standards contained between 10 and 1000 ppm alpha-amylase.

When needed, water and solvent concentrations were established by capillary gas chromatography. A Hewlett-Packard 6890 gas chromatograph equipped with capillary inlets, thermal conductivity detectors (TCD), HP-7683 autoinjectors, and a Chemstation was used for these analyses. Samples and calibration standards were diluted 5:1 with tetrahydrofuran (THF), which was used as the diluent, internal standard, and reference solvent. Calibrations were made using a 50 weight percent solvent in water solution after establishing the linearity range for the analyses. Data were recorded as the average of multiple injections.

Examples 1 to 11 and Comparative Examples C-1 and C-2Single-Liquid System:

5 The whole fermentation broth containing α -amylase BD5088 (a heat stable, low pH enzyme with a molecular weight of ~49 kDaltons (kDa) and an isoelectric point (pI) of 4.5), obtained by fermentation of the production organism *Pseudomonas fluorescens*, was heat treated for 30 minutes at 70 °C. An organic solvent that is miscible in water was added to the fermentation broth. After agitation of the mixture for about 30
10 minutes, only one liquid phase is present. In most cases, biomass solids were separated from the resulting liquid using a high-speed batch centrifuge. The results are shown in Table 1 that follows. A number of the examples listed in Tables 1 to 3 are duplicate experiments that were carried out to better understand experimental variability. Although the results show some variability, the trends in the data are consistent.

15 Two types of examples were performed. One set of examples involved adding 15 grams of the solvent to 15 grams of the fermentation broth. These examples are indicated in Table 1 by the solvent to fermentation broth ratio of 1.0. The other set of examples involved first diluting 7.5 ml of the fermentation broth with 7.5 ml of water
20 and then adding 15 ml of the solvent. This resulted in a significantly higher ratio of extraction liquid to the fermentation broth, although the concentration of the solvent in the liquid after extraction was not greatly different. These examples are indicated in Table 1 by the solvent to fermentation broth ratio between 2.6. In this case, this ratio is defined as the ratio of the weight of added liquid, including water and solvent to the
25 weight of the initial feed broth, prior to extraction.

Table 1. Results of Whole Broth Extractions Involving a Single Liquid Phase

Example	Solvent System	Temp	Solvent to Feed Ratio Weight of Added Liquid to Weight of Feed Broth prior to Extraction	Concentration of Enzyme in the Liquid after Extraction	Recovery of Enzyme from the Feed Broth into the Liquid after Extraction
		(°C)	(wt / wt)	(ppm-wt)	(%)
C-1*	Water alone	22	1.0**	95	3
C-2*	Water alone	50	1.0**	178	6
1	IPA-water	22	1.0**	237	7
2	IPA-water	22	1.0**	199	6
3	IPA-water	22	1.0**	350	11
4	IPA-water	22	2.6***	638	35
5	EB-water	22	1.0**	908	27
6	EB-water	22	1.0**	1900	57
7	PM-water	22	1.0**	2920	88
8	PM-water	22	1.0**	2850	86
9	PM-water	22	1.0**	2690	81
10	PM-water	22	1.0**	2780	84
11	PM-water	50	1.0**	3300	99

* not an example of the present invention.

** equal weights of whole broth and solvent were mixed.

5 *** diluted 7.5 ml broth with 7.5 ml of water, then added 15 ml of solvent.

The concentration of enzyme in the feed broth was approximately 6380 ppm. The concentration of biomass was 8.7 wt%.

10 In all of the examples in Table 1, broth that had been stored frozen for several months was used.

The liquid-phase enzyme concentrations determined by high pressure liquid chromatography (HPLC) analysis were used to calculate the amount of enzyme recovered from the biomass solids into the liquid phase. Since the enzyme is expressed

as inclusion bodies within the cell, it is originally associated with the biomass. The data in Table 1 show that a significant fraction of the α -amylase present as inclusion bodies within the cells are efficiently extracted out of a heat treated whole fermentation broth into a single liquid phase containing roughly equal amounts of water and glycol ether without pretreatment steps specifically designed to lyse the cells.

The results of Yes/No enzyme activity tests performed on the extracted liquid are shown in Table 2 that follows.

Table 2. Yes/No Activity Test Results for Examples Shown in Table 1

Example	Solvent System	Temp (°C)	Measured Enzyme Concentration in the Extract Liquor	Concentration of Enzyme in the Activity Test Sample after Sample Preparation (ppm)	Measured Absorbance	Absorbance Expected for that Enzyme Concentration Assuming No Loss of Bioactivity	Activity Retained? (Yes/No)
1	IPA-water	22	237	0.4	3.3	1.5+	Yes
5	EB-water	22	908	1.3	3.02	2.9+	Yes
C-1	Water alone	22	95	0.16	3.53	0.5+	Yes
C-2	Water alone	50	178	0.28	2.37	1.0+	Yes

Examples 12 to 19Two-Liquid System

5 Equal weights of glycol ether and a whole fermentation broth were mixed at a temperature that allowed formation of two phases, that is, an aqueous liquid phase and an organic liquid phase rich in the glycol ether. The α -amylase enzyme recovery was increased roughly to 50% for EB (at 60 °C), and 70% for PnP (at 22 °C). Although the organic phase contained up to 33% dissolved water by weight, less than 10 parts per million (ppm) of enzyme was detected in the organic phase. Essentially all of extracted
10 α -amylase enzyme was present in the liquid aqueous phase. The results obtained are given in Tables 3 and 4 that follow. A number of the examples given in Tables 3 and 4 are duplicate experiments that were carried out to better understand experimental variability. Although the results show some variability, the trends in the data are consistent.

Table 3. Results of Whole Broth Extractions Involving Two Liquid Phases

Example	Solvent System	Temp	Weight of Added Solvent to Weight of Feed Broth prior to Extraction	Concentration of Enzyme in the Aqueous Phase after Extraction**	Concentration of Glycol Ether in the Aqueous Phase	Concentration of Water in the Organic Phase
		(°C)	(wt / wt)	(ppm-wt)	(wt %)	(wt %)
12	PnP-water	22	1.0	5350	22	33
13	PnP-water	22	1.0	5180	22	33
14	EB-water	60	1.0	3060	17	31
15	EB-water	60	1.0	3750	17	31
16	PnB-water	50	1.0	250	4.9	12.8
17	PnB-water	22	1.0	190	5.6	13.3
18	PnB-water	22	1.0	160	5.6	13.3
19	PnB-water	-5	1.0	177	5.0	15.4

Table 4. Results of Whole Broth Extractions Involving Two Liquid Phases

Example	Solvent System	Temp	Weight of Feed Broth	Approx. Weight of Aqueous Phase after Extraction*	Approx. Weight of Organic Phase after Extraction*	Recovery of Enzyme from the Feed Broth into the Aqueous Phase after Extraction**
		(°C)		(grams)	(grams)	(%)
12	PnP-water	22	15	13	17	70
13	PnP-water	22	15	13	17	70
14	EB-water	60	15	13	17	40
15	EB-water	60	15	13	17	50
16	PnB-water	50	10	9	11	4
17	PnB-water	22	10	9	11	3
18	PnB-water	22	10	9	11	2
19	PnB-water	-5	10	9	11	3

* determined from material balance

** less than 10 ppm enzyme was detected in the organic phase after extraction

- 5 The concentration of enzyme in the feed broth was approximately 6380 ppm. The concentration of biomass was 8.7 wt%.

The results of Yes/No enzyme activity tests performed on the extracted liquid are shown in Table 5 that follows.

Table 5. Yes/No Activity Test Results for Examples given in Tales 3 and 4

Sample	Solvent System	Temp	Measured Enzyme Concentration in the Extract Liquor	Concentration of Enzyme in the Activity Test Sample after Sample Preparation	Measured Absorbance	Absorbance Expected for that Enzyme Concentration Assuming no Loss of Bioactivity	Activity Retained Yes/No
15	EB-water						
	Aqueous Layer	60 °C	3750	197	3.0	3.7+	Yes, although some loss of activity is apparent
	Organic Layer	60 °C	14	0.7	2.5	1.5+	Some activity is detected, even though only traces of enzyme were present in the sample
13	PnP-water						
	Aqueous Layer	22 °C	5180	No data*	----	----	No data*
	Organic Layer	22 °C	15	0.8	0.2	1.5+	Little activity

* the aqueous phase was too small to test activity

The results of enzyme activity tests performed on the extracted liquid using a test utilizing *p*-nitrophenyl- α -D-hexa-glucopyranoside are shown in Table 6 that follows.

5 Table 6. *p*-Nitrophenyl- α -D-hexa-glucopyranoside Activity Test Results for Examples given in Tales 1, 3 and 4

Solvent	Experiment No.	Temp (°C)	Concentration of Enzyme in Extract Liquor (aqueous phase)		Activity Retained (%)
			LC (ppm)	Activity Assay (ppm)	
11	PM	50	3300	2900	88
10	PM	22	2800	2800	100
12	PnP	22	5370	4570	85
6	EB	22	1900	1040	55
14	EB	60	3060	2470	80
3	IPA	22	350	300	86
16	PnB	50	250	255	100
18	PnB	22	160	155	97

Experiments were carried out to determine if enzyme already dissolved in water would transfer into a separate glycol ether phase. The solvents used in these experiments were ethylene glycol n-butyl ether (EB), propylene glycol n-propyl ether (PnP), dipropylene glycol n-propyl ether (DPnP), tripropylene glycol n-propyl ether (TPnP), and dipropylene glycol dimethyl ether (DMM). The enzyme feed solutions were prepared from a purified α -amylase enzyme. Equal weights of the aqueous enzyme feed and the added solvent were used to maximize the potential for transfer of the enzyme there. In certain cases, this required operating at temperatures slightly above the LCST. In the case of EB, it required operating at 70 °C. In all of these examples, only trace amounts of the enzyme were observed to transfer from the aqueous phase into the glycol ether phase. This was true even when significant amounts of water were present in the glycol ether phase (up to about 52 wt%). Adjusting the pH over the range of 4-7 using buffer solutions did not alter this behavior. Using a pH of 2 caused precipitation of the enzyme but did not result in transfer of significant amounts of the enzyme into the glycol ether phase. The results obtained are given in Table 7 that follows.

Table 7. Results of Extraction Experiments Using Purified Enzyme Solution

Experiment #	Experimental Conditions				Aqueous Layer (wt.%)			Organic Layer (wt.%)			Activity Test (Absorbance)	
	Glycol Ether	ppm enzyme in aq. Feed	pH of Enzyme Solution	Sampling Temp (C)	Glycol Ether	Water	Amylase (ppm)	Glycol Ether	Water	Amylase (ppm)	Aqueous Layer	Organic Layer
1	PnP	345.3	4	38	26.26	73.68	675	59.78	40.22	0.0	3.55	3.35
2	PnP	329.4	4	34	29.5	70.43	723	57.16	42.84	0.0	3.44	1.87
3	PnP	334.2	2	32	23.09	76.91	0.0	62.47	37.53	10.4	0.28	N/A*
4	DPnP	335.7	7	25	15.96	84.00	374	78.5	21.50	0.0	3.63	0.43
5	DPnP	345.3	4	17	30.99	68.97	400	62.89	37.11	0.0	3.72	0.93
6	DMM	334.6	5	10	42.51	57.47	182	94.93	5.07	0.0	3.40	0.09
7	DMM	335.9	4	10	43.96	56.02	187	94.19	5.81	0.0	3.46	2.02
8	TPnP	334.6	5	10	25.91	74.06	306	77.95	22.05	0.0	3.32	0.70
9	TPnP	335.9	4	10	31.08	68.89	338	76.92	25.86	0.0	3.46	0.03
10	EB	235.4	4	70	15.57	84.4	262	48.86	51.13	11.8	3.36	1.64

*: α -amylase precipitated out of solution at this pH

20

The activity test results given in Table 7 indicate full activity for all of the aqueous samples. These solutions contained the enzyme plus a considerable amount of dissolved solvent. Interestingly, significant activity was also measured for some of the organic samples, even though LC analysis of the organic layer did not detect the presence of the enzyme. These results provide another example of how even very small

25

amounts of the enzyme (< 1 ppm in the test solutions) can catalyze significant hydrolysis of starch in the test.

5 The data in Table 7 also show that the enzyme solution may be concentrated to some extent by extraction of water into a pre-dried glycol ether phase; that is, one that is unsaturated with water prior to its use in the extraction. For example, experiments 1 and 2 demonstrate that PnP can double the α -amylase concentration by extracting water from the aqueous layer and reducing the size of the aqueous layer in half. Experiments 4, 5, and 10 demonstrate that DPnP and EB also yield a concentration effect. The TPnP and DMM, on the other hand, showed little if any concentration effect because an equal
10 amount of glycol ether (approximately) also transferred into the aqueous phase.

 Because the enzyme is retained almost exclusively in the aqueous phase, transfer of water and organic impurities out of the aqueous phase into the glycol ether
15 phase can provide an opportunity to purify and concentrate an aqueous enzyme solution.

 Other embodiments of the invention will be apparent to those skilled in the art from a consideration of this specification or practice of the invention disclosed herein.
20 It is intended that the specification and examples be considered as exemplary only, with the true scope and spirit of the invention being indicated by the following claims.

WHAT IS CLAIMED IS:

1. A method for extracting an intracellular protein from a fermentation broth comprising the steps of:

5

(a) intermixing a sufficient quantity of a water miscible alcohol or glycol ether with an aqueous fermentation broth at a temperature at which the alcohol or glycol ether forms a single aqueous solution comprising a protein, the water miscible alcohol or glycol ether, and water;

10

(c) separating the aqueous phase comprising the protein, the water miscible alcohol or glycol ether, and water formed in step (a) from solid biomass impurities; and, optionally,

15

(c) recovering the protein from the aqueous phase obtained in step (b) by any conventional protein recovery method.

2. The method of Claim 1, wherein the alcohol is methanol, ethanol, 2-propanol, or 2-methyl-2-propanol.

20

3. The method of Claim 2, wherein the alcohol is 2-propanol

4. The method of Claim 1, wherein the glycol ether is completely miscible with water in the temperature range of from about 20 °C to about 80 °C.

25

5. The method of Claim 4, wherein the glycol ether is ethylene glycol n-propyl ether, propylene glycol ethyl ether, propylene glycol methyl ether, diethylene glycol n-butyl ether), diethylene glycol ethyl ether, diethylene glycol methyl ether, triethylene glycol n-butyl ether, triethylene glycol n-pentyl ether, triethylene glycol ethyl ether, triethylene glycol methyl ether, or diethylene glycol dimethyl ether.

30

6. The method of Claim 1, wherein the glycol ether is miscible with water at the temperature of about 20 °C but become immiscible as the water-ethylene glycol solution is heated.

35

7. The method of Claim 6, wherein the glycol ether is ethylene glycol n-butyl ether, ethylene glycol iso-butyl ether, propylene glycol n-propyl ether, dipropylene glycol ethyl ether, dipropylene glycol iso-propyl ether, diethylene glycol 2-methylbutyl ether, diethylene glycol n-pentyl ether), triethylene glycol n-heptyl ether, triethylene glycol n-hexyl ether, diethylene glycol ethyl ether acetate, or diethylene glycol diethyl ether.
8. A method for extracting an intracellular protein from a fermentation broth comprising the steps of:
- (d) intermixing a sufficient quantity of a partially water miscible glycol ether with an aqueous fermentation broth at a temperature where the glycol ether forms an aqueous phase comprising a protein, partially water miscible glycol ether, and water; and an organic phase comprised mainly of partially miscible glycol ether;
 - (e) separating the liquid organic phase formed in step (a) from the aqueous phase,
 - (e) separating the aqueous phase obtained in step (b) from solid biomass impurities; and, optionally,
 - (d) recovering the protein from the aqueous phase obtained in step (c) by any conventional protein recovery method.
9. The method of Claim 8, wherein the glycol ether is miscible with water at the temperature of about 20 °C but become immiscible as the water-ethylene glycol solution is heated.
10. The method of Claim 9, wherein the glycol ether is ethylene glycol n-butyl ether, ethylene glycol iso-butyl ether, propylene glycol n-propyl ether, dipropylene glycol ethyl ether, dipropylene glycol iso-propyl ether, diethylene glycol 2-methylbutyl ether, diethylene glycol n-pentyl ether), triethylene glycol n-heptyl ether, triethylene glycol n-hexyl ether, diethylene glycol ethyl ether acetate, or diethylene glycol diethyl ether.

11. The method of Claim 8, wherein the glycol ether forms a separate phase with water at about 20 °C and separate further upon heating.

12. The method of Claim 11, wherein the glycol ether is ethylene glycol 2-
5 methylbutyl ether, ethylene glycol n-hexyl ether, ethylene glycol n-pentyl ether,
propylene glycol n-butyl ether, propylene glycol tert-butyl ether, propylene glycol iso-
propyl ether, dipropylene glycol n-butyl ether, dipropylene glycol n-propyl ether,
diethylene glycol n-hexyl ether, tripropylene glycol n-butyl ether, tripropylene glycol n-
propyl ether, ethylene glycol ethyl ether acetate, ethylene glycol n-butyl ether acetate,
10 diethylene glycol n-butyl ether acetate, propylene glycol methyl ether acetate, ethylene
glycol diethyl ether, ethylene glycol dibutyl ether, diethylene glycol dibutyl ether, or
dipropylene glycol dimethyl ether.

ABSTRACT

A method for extracting an intracellular enzyme from a whole fermentation
broth using a water miscible alcohol, or a water miscible or partially water miscible
5 glycol ether.